

## FORMULATION

Formulation No. 170607E-2-J1

PHASE	INCI NAME	COMMERCIAL NAME	Q'TY
A	Water	Water	30.70
A	Glycerin	Glycerin	5.00
A	3-O-Ethyl Ascorbic Acid	Et-VC™	2.00
A	Magnesium Sulfate	MgSO <sub>4</sub> · 7H <sub>2</sub> O	0.50
A	Sodium Chloride	NaCl	0.80
B	Cyclopentasiloxane (And) Titanium Dioxide (And) Cyclohexasiloxane (And) Iron Oxides (C.I. 77492) (And) Iron Oxides (C.I. 77491) (And) PEG/PPG-18/18 Dimethicone (And) Iron Oxides (C.I. 77499) (And) Dimethicone/Vinyl Dimethicone Crosspolymer (And) Disteardimonium Hectorite (And) Tocopheryl Acetate (And) Isopropyl Titanium Triisostearate (And) Triethoxysilylethyl Polydimethylsiloxyethyl Dimethicone	KOBOgel D5 light	32.00
B	Cetyl PEG/PPG-10/1 Dimethicone	Jeesilc EM90	2.50
B	Cyclopentasiloxane (and) Caprylyl Dimethicone Ethoxy Glucoside	BELSIL SPG 128 VP	5.00
B	Caprylyl Glycol (and) Phenoxyethanol (and) Hexylene Glycol	Jeecide CAP-2	1.00
B	Polymethyl Methacrylate	Ganzpearl GM 0600	2.00
B	Silica	MSS-500W	2.50
B	Diisostearyl Malate	Corum 5015	2.00
B	Trimethylsiloxy silicate	BELSIL TMS 803	1.00
B	Dimethicone	BELSIL DM 5	5.00
C	Cyclopentasiloxane (and) Cyclohexasiloxane (and) Quaternium-90 Bentonite (and) Propylene Carbonate	Tixogel-VSP 1438	5.00
C	Cyclopentasiloxane (and) Dimethicone/Vinyl Dimethicone Crosspolymer	BELSIL EG 2	3.00

### INSTRUCTIONS

1. Mix phase A uniformly.
2. Mix phase B uniformly.
3. Mix phase C uniformly.
4. Add phase A into phase B, then mix uniformly.
5. Add phase C into phase A/B, then mix uniformly.

### APPEARANCE

Beige Cream

### TECHNICAL DATA

pH (Cyber Scan pH510): N/A

Viscosity (Brookfield DV-1I/S64, 3 rpm, 30 sec): 130000 cP